

ENGLISH

S-2288™

For Laboratory Use Only

S-2288™

S-2288 is a chromogenic substrate sensitive to a broad spectrum of serine proteases.

COMPOSITION

Each vial contains chromogenic substrate S-2288, 25 mg and mannitol 40 mg as a bulking agent.

CHEMISTRY

Chemical name: H-D-Isoleucyl-L-prolyl-L-arginine-p-nitroaniline dihydrochloride.

Formula: H-D-Ile-Pro-Arg-pNA · 2HCl

Mol. wt: 577.6

$\epsilon_{316 \text{ nm}^1}$: $1.27 \cdot 10^4 \text{ mol}^{-1} \cdot \text{L} \cdot \text{cm}^{-1}$

Solubility: > 40 mmol/L in H₂O

Stability: Substance: Stable until expiry date if stored at 2-8°C. Avoid exposure to light. The substance is hygroscopic and should be stored in a dry place.

Solution: 10 mmol/L in H₂O is stable for more than two months at 2-8°C. Contamination by microorganisms may cause hydrolysis.

Suitable

stock solution: 10 mmol/L in H₂O

KINETIC DATA

The following kinetic constants were estimated at 37°C in Tris buffer pH 8.4, I 0.15.

Enzyme	K _m (mol/L)	V _{max} (mol/min and unit)	k _{cat} (sec ⁻¹)
Thrombin	$3 \cdot 10^{-6}$	$10 \cdot 10^{-8}$	118
Urokinase	$2 \cdot 10^{-4}$	$2 \cdot 10^{-10}$	16
Factor XII f	$4 \cdot 10^{-4}$	$5 \cdot 10^{-5}$	23
t-PA ¹¹ one-chain	$1 \cdot 10^{-3}$	$10 \cdot 10^{-11}$	26
t-PA ¹¹ two-chain	$3 \cdot 10^{-4}$	$11 \cdot 10^{-11}$	28
Plasma kallikrein	$1 \cdot 10^{-3}$	$13 \cdot 10^{-7}$	-
Plasmin	$9 \cdot 10^{-3}$	$5 \cdot 10^{-6}$	181
Factor Xa	$2 \cdot 10^{-3}$	$9 \cdot 10^{-6}$	110
C _s	$3 \cdot 10^{-3}$	$3 \cdot 10^{-6}$	4
C _{tr}	$6 \cdot 10^{-4}$	$1 \cdot 10^{-6}$	2

¹¹ Tissue plasminogen activator (porcine)

Units expressed in nmols are as follows:

Thrombin (human) Ortho	1 NIH U	= $1.4 \cdot 10^{-2}$ nmol
Urokinase Leo	1 Ploug U	= $1.8 \cdot 10^{-4}$ nmol
Factor XIII* Mol wt 28 000	1 mg	= 36 nmol
t-PA (one- and two-chain)		
Mol wt 64 000	1 IU	= $6.3 \cdot 10^{-6}$ nmol(1)
Plasma kallikrein	U	= $\mu\text{mol}/\text{min}$
Plasmin (human)Chromogenix	1 CU	= 0.45 nmol
Factor Xa (bovine) Diagen	1 U (Denson)	= $1.4 \cdot 10^{-2}$ nmol
C _s * and C _{tr} * Mol wt 85 000	1 mg	= 12 nmol

*The enzyme is assumed to be pure

STANDARDIZATION

With a substrate concentration of $1 \cdot 10^{-3}$ mol/L and an enzyme concentration of $4 \cdot 10^{-9}$ mol/L the following activities are obtained.

Enzyme	$\Delta A/\text{min}$	Enzyme	$\Delta A/\text{min}$	Enzyme	$\Delta A/\text{min}$
Thrombin	0.275	Plasmin	0.042	Factor Xa	0.084
Urokinase	0.031	t-PA (one-chain)	0.030	C _s	0.002
Factor XIII	0.023	t-PA (two-chain)	0.040	C _{tr}	0.003

The tissue plasminogen activator is not affected by the following inhibitors used at the concentrations given.

1. Trasylol 30 KIU/mL.
2. Soybean trypsin inhibitor 50 $\mu\text{g}/\text{mL}$.
3. Antithrombin 0.1 PEU/mL and heparin 3 IU/mL.

APPLICATIONS

The substrate has been used for the determination of Tissue plasminogen activator in purified preparations (2,3,4)

DEUTSCH

S-2288™

Nur für Laborzwecke

S-2288 ist ein, auf ein breites Spektrum Serinproteasen sensitives, chromogenes Substrat.

ZUSAMMENSETZUNG

Jedes Fläschchen enthält 25 mg chromogenes Substrat S-2288 und 40 mg Mannitol als Füllstoff.

CHEMIE

Chemischer Name: H-D-Isoleucyl-L-Prolyl-L-Arginin-p-Nitroanilin Dihydrochlorid

Formel: H-D-Ile-Pro-Arg-pNA · 2HCl

Molekulargewicht: 577,6

$\epsilon_{316 \text{ nm}^1}$: $1,27 \cdot 10^4 \text{ mol}^{-1} \cdot \text{L} \cdot \text{cm}^{-1}$

CHROMOGENIX

Chromogenix-
Instrumentation Laboratory SpA
V.le Monza 338 - 20128 Milano (Italy)

301959RO



REFERENCES

1. WALLÉN P RÅNBY M: Personal communication.
2. WALLÉN P RÅNBY M: Peptides of the Biological Fluids. Brussels, Belgium. Abstr. 67, (1980)
3. FRIBERGER P et al: Progr. in Chem. Fibr. and Thromb. IV Ed JF Davidson. 149 (1979)
4. FRIBERGER P: Thesis Scand J clin lab Invest 42 suppl 162, 58-59 (1982)
5. Chromogenix AB: Determination of Tissue Plasminogen Activator (t-PA) in purified preparations. Laboratory instructions.
6. Chromogenix AB: Determination of proteolytic activity in plasma, serum or Euglobulin fractions. Laboratory instructions.

CHROMOGENIX